

How large is the contribution of cosmic web to Ω_Λ ?

A preliminary study on a novel inhomogeneous model

Stefano Viaggiu

Dipartimento di Matematica, Università "Tor Vergata" Roma,
Via della Ricerca Scientifica, 1
Rome, Italy 00133,
viaggiu@axp.mat.uniroma2.it

Marco Montuori

SMC-ISC-CNR and Dipartimento di Fisica,
Università "La Sapienza" Roma,
Ple. Aldo Moro 2
00185, Rome, Italy,
marco.montuori@roma1.infn.it,

May 23, 2013

Abstract

The distribution of matter in the universe shows a complex pattern, formed by cluster of galaxies, voids and filaments denoted as cosmic web. Different approaches have been proposed to model such structure in the framework of the general relativity. Recently, one of us has proposed a generalization (Λ FB model) of the Fractal Bubble model, proposed by Wiltshire, which accounts for such large scale structure. The Λ FB model is an evolution of FB model and includes in a consistent way a description of inhomogeneous matter distribution and a Λ term. Here we analyse the Λ FB model focusing on the relation between cosmological parameters. The main result is the consistency of Λ CDM model values for $\Omega_{\Lambda 0}$ (≈ 0.7) and Ω_{k0} ($|\Omega_{k0}| < \approx 0.01$) with a large fraction of voids. This allows to quantify to which extent the inhomogeneous structure could account for Λ constant consistently with standard values of the other cosmological parameters.

PACS Numbers: 98.80.-k, 98.80.Jk, 95.36.+x, 04.20.-q

1 Introduction

The model of universe which currently gives the best fit of the available astrophysical observations is the Λ Cold Dark Matter model (Λ CDM). The

model is based on an exact solution of the equation of General Relativity which assumes homogeneity and isotropy with a F-L metric. Quite soon, however, observational evidences required the addition of two main artefacts to the theory: the dark matter and the dark energy. Rotation curve of spiral galaxies, velocity dispersion of galaxies in galaxy clusters, cluster mass estimate from hot intra-cluster gas emission, gravitational lensing from galaxy cluster, large scale structure formation from CMB tiny fluctuations required an addition of dark matter. The observation of an accelerated expansion of the universe required the addition of the dark energy. According to the best fit model, dark matter should account for $\approx 23\%$ of the total energy density, while the dark energy for $\approx 73\%$ ([1, 2, 3]). Notwithstanding, the nature of both remains unknown and is amongst the deepest problems of modern physics. For these reasons in the past decade, several authors [4]-[29] have explored alternatives to the Λ CDM model.

The discovery of a lumpier universe than expected offered a possibility in this respect. The universe appears indeed organized as a cosmic web, which can be described as an interconnected network: spherical clusters form the nodes and are joined by elongated filaments defining 2D sheets. Recent analysis have shown that voids in the network could fill between $\approx 40\% - 70\%$ of the space (accordingly to definition and measure of the voids) and have a continuous distribution of sizes depending on the galaxy sample selected [30, 31, 32, 33]. Such observed inhomogeneities are usually described as a first order perturbation with respect to the homogeneous and isotropic exact solution. This formulation has a limited application, since it is valid for small perturbation $\delta\rho/\rho$ and cannot describe the lumpy universe at low redshift. Nevertheless, it was soon clear, mainly through the formulation of the Buchert formalism, that such an inhomogeneous matter distribution could indeed mimic the presence of dark energy [34, 35, 36, 37, 38]. Since then, some authors explored this possibility to the extreme consequence to explain the whole amount of dark energy on the basis of inhomogeneous distribution. Our personal opinion is that the requested existence of Gpc scale voids makes these models questionable. From the other side, the standard perturbative approach of Λ CDM cannot answer to the question of the effect of small scale strong fluctuations on larger scale. In this context, an interesting line appears the exploration of cosmological models including the largely accepted cosmological constant paradigm and at the same time a non perturbative approach to the observed inhomogeneities. This issue is a very recent one (see [39]) and so far limited to the not so realistic (even useful) case of spherical symmetry. Going beyond such studies and toward a more realistic model of matter distribution in the universe is the target of the present paper. The elaboration of a similar model will allow, for example, to study in a non perturbative way the percentage of dark energy that can be explained in terms of the observed inhomogeneities or the role of voids in the process of structure formation (see for example [40]) in a more complete

way for the presence both of dark energy and voids in the model. More in general, it would be possible to explore the effect of the observed inhomogeneities onto cosmic scale data, as barionic acoustic oscillations and cosmic radiation.

Our baseline is the so-called 'fractal bubble' FB or 'timescape' cosmology [22, 23]. It is based on the Buchert average scheme [18] and introduces a non-uniform time flow with two scales corresponding to the voids and walls of a schematic lattice cosmic structure. In the FB model the apparent acceleration is explained in terms of different rate of clocks located in walls (our point of observation) as compared to the clocks of voids. The viability of the FB or Timescape model as an alternative to the Λ CDM model is still matter of debat [24, 25]. In any case, the FB model is the first one which describes a schematic cosmic web in a non perturbative way and without invoking particular symmetries. This interesting feature has prompted one of us to propose an inhomogeneous model based on the same partitioning within the Buchert average scheme, but including a $\Lambda \neq 0$ and an uniform time flow [41]. This model, hereafter Λ FB model, allows to investigate, in a non perturbative way, the effects of large scale spatial inhomogeneities and non-vanishing curvature on the cosmological parameters. The present preliminary study is devoted to such aim. The scheme of the paper is the following: in section 2 we write the relevant equations of the model. We obtain a new interesting equation involving all the parameters of the model evaluated at the present epoch, which is analysed in sections 3 and 4. In section 5 we 'dress' the cosmological parameters and finally, we report our conclusions in Section 6.

2 Basic equations

The Λ FB model [41] incorporates the observed present inhomogeneities within the cosmological constant paradigm. The starting point is the FB model introduced by Wiltshire [22, 23]. This model considers a matter distribution in the universe as a regular network formed by walls and voids. The Λ FB model assumes the same partitioning of the two scale FB model [23]: regions called 'finite infinity' (F_I) which are the set of timelike boundaries of compact **disjoint** domains I , with a zero average expansion and a positive one outside, i.e. $\langle \theta \rangle_{\cup_I F_I} = 0$. The F_I regions are within 'wall regions' whose metric is, on average:

$$ds_w^2 = -dt^2 + a_w^2 [d\eta_w^2 + \eta_w^2 d\Omega^2]. \quad (1)$$

Our position is in F_I . The regions complementary to the walls with respect to the particle horizon are called voids and are expanding with an average hyperbolic metric given by:

$$ds_v^2 = -dt^2 + a_v^2 [d\eta_v^2 + \sinh^2(\eta_v) d\Omega^2]. \quad (2)$$

Note that, contrary to the FB model, in our model the time flow is isotropic. This is due to the presence of a non vanishing cosmological constant, which is absent in the FB model since it intends to explain it as an effect of inhomogeneity. The isotropic flow has the useful feature to avoid the shortcomings related to the junction conditions of the original FB model (see [42]). In any case, a nontrivial phenomenological lapse function involves new physics relating to gravitational energy which is not yet widely accepted, and we will therefore consider the commonly accepted alternative that time flows uniformly.

The Hubble parameters in walls and voids are respectively $H_w = \frac{\langle \theta \rangle_w}{3} = \frac{\dot{a}_w}{a_w}$, $H_v = \frac{\langle \theta \rangle_v}{3} = \frac{\dot{a}_v}{a_v}$. An important assumption of the Λ FB model (and FB one) is the existence of a scale of homogeneity with the scale factor $a(t)$. The average on the whole volume of the particle horizon $V = a^3 V_i$ is:

$$a^3 = f_{wi} a_w^3 + f_{vi} a_v^3, \quad f_{wi} + f_{vi} = 1, \quad (3)$$

where f_{wi} and f_{vi} are the fractions of walls and voids at the time $t = t_i$. We choose the time t_i (the initial time) as the recombination time in order to compare the model with available observational data. In general the following relations are valid for any time:

$$f_v(t) + f_w(t) = 1, \quad f_w = f_{wi} \frac{a_w^3}{a^3}, \quad f_v = f_{vi} \frac{a_v^3}{a^3}. \quad (4)$$

The Hubble rate H at the scale of homogeneity is:

$$H = f_w H_w + f_v H_v, \quad H = I_w H_w = I_v H_v. \quad (5)$$

and the density parameters are:

$$\Omega_m = \frac{8\pi \langle \rho \rangle}{3H^2}, \quad \Omega_k = -\frac{\langle \mathcal{R} \rangle}{6H^2}, \quad \Omega_{\mathcal{Q}} = -\frac{\mathcal{Q}}{6H^2}, \quad \Omega_{\Lambda} = \frac{\Lambda}{3H^2}, \quad (6)$$

where $\mathcal{Q}$ is the kinematic backreaction and $\mathcal{R}$ is the spatial curvature. The Buchert equations are:

$$\Omega_m + \Omega_{\Lambda} + \Omega_k + \Omega_{\mathcal{Q}} = 1, \quad (6\mathcal{Q} + 2\langle \mathcal{R} \rangle)\dot{a} + a[\dot{\mathcal{Q}} + \langle \dot{\mathcal{R}} \rangle] = 0. \quad (7)$$

Equation (7) can be easily manipulated to obtain the final equations for the model (for more details see [41]):

$$(1 - f_v) \frac{\dot{a}}{a} - \frac{\dot{f}_v}{3} = \sqrt{\Omega_{m0} H_0^2 \frac{a_0^3}{a^3} \left(\frac{1 - \epsilon_i}{\Omega_F} \right) (1 - f_v)}, \quad (8)$$

$$\frac{\dot{a}}{a} + \frac{\dot{f}_v}{3f_v} = \frac{a_0 H_0}{a f_v^{\frac{1}{3}}} \sqrt{\frac{\Omega_{k0}}{f_{v0}^{\frac{1}{3}}} + \frac{\Omega_{\Lambda 0} a^2}{f_v^{\frac{1}{3}} a_0^2} + \frac{a_0 \Omega_{m0}}{a f_v^{\frac{1}{3}}} \left(1 + \frac{\epsilon_i - 1}{\Omega_F} \right)}, \quad (9)$$

where ϵ_i is a integration constant. A first integral of eqs (8, 9) is:

$$\frac{(1 - \epsilon_i)I_w^2 \Omega_m}{(1 - f_v)} = \Omega_F = e^{-\int_{t_i}^t \frac{\Lambda}{H_w} dt}. \quad (10)$$

From equation (8) (see [41]) we can solve for a_w and Ω_F :

$$a_w = a_{w0} \sinh^{\frac{2}{3}} \left(\frac{\sqrt{3\Lambda}}{2} t \right), \quad \Omega_F = \left(\frac{\cosh \left(\sqrt{\frac{3\Lambda}{4}} t_i \right)}{\cosh \left(\sqrt{\frac{3\Lambda}{4}} t \right)} \right)^2. \quad (11)$$

We are now able to derive an useful and interesting formula. By combining equations (8) and (9) and evaluating them at the present epoch t_0 we obtain:

$$1 = \sqrt{\Omega_{m0}(1 - f_{v0}) \frac{(1 - \epsilon_i)}{\Omega_{F0}}} + \sqrt{f_{v0}} \sqrt{\Omega_{k0} + \Omega_{\Lambda 0} + \Omega_{m0} \left(1 + \frac{\epsilon_i - 1}{\Omega_{F0}} \right)}, \quad (12)$$

The equation (12) constraints the density parameters and the fraction of voids at the present time t_0 . The equation was already present in [41]. Now we intend to go a step beyond, expressing the constant Ω_{F0} in terms of density parameters. Integrating equation (8) we get:

$$(1 - f_v)^{\frac{1}{3}} a = a_0 \left(\frac{\Omega_{m0}(1 - \epsilon_i)}{\Omega_{\Lambda 0}} \right)^{\frac{1}{3}} \frac{\sinh^{\frac{2}{3}} \left(\frac{\sqrt{3\Lambda}}{2} t \right)}{\cosh^{\frac{2}{3}} \left(\frac{\sqrt{3\Lambda}}{2} t_i \right)}. \quad (13)$$

It should be noticed that in the equation (13), according to [22] and without loss of generality, we have set to zero the integration constant. In such a way, the study of the cosmological constant is simpler. This choice constraints the integration constant in the integration of the equation (9) (see [22]). Moreover, since the current estimation of Λ is very small and $t_i \approx 3.77 \cdot 10^{-5}$ Gyr, we can set with good approximation $\cosh^{\frac{2}{3}} \left(\frac{\sqrt{3\Lambda}}{2} t_i \right) \simeq 1$. Evaluating equation (13) at the present epoch, we get:

$$\frac{(1 - \epsilon_i)}{\Omega_{F0}} = 1 - \epsilon_i + \frac{\Omega_{\Lambda 0}}{\Omega_{m0}} (1 - f_{v0}). \quad (14)$$

and by putting (14) in (12) we have:

$$1 = \sqrt{(1 - f_{v0})[\Omega_{m0}(1 - \epsilon_i) + \Omega_{\Lambda 0}(1 - f_{v0})]} + \sqrt{f_{v0}} \sqrt{\Omega_{k0} + \Omega_{m0}\epsilon_i + f_{v0}\Omega_{\Lambda 0}}. \quad (15)$$

The formula (15) was not present in [41]; it involves all the parameters of the Λ FB model and has been obtained by using all the relevant equations

(i.e. (8) and (9)). Moreover, it gives a compatibility equation for cosmological parameters that are averaged quantities in principle measurable. It should be noted that this equation is not perturbative. It allows to study the effects of the inhomogeneities without ruling out the dark energy and avoiding the shortcomings of the perturbation theory when applied to large inhomogeneities.

The cosmological parameters present in (15) are different from those of the Λ CDM model. The concordance model is based on an exact solution of Einstein's equations and the corresponding cosmological parameters are local quantities. The homogeneity and isotropy of the solution implies that the cosmological parameters are the same in any spatial point for any fixed comoving time t . As the timescape model, also the Λ FB one is obtained within the Buchert formalism, where the cosmological parameters are averaged non local quantities. In this formalism for a given scalar ψ at a fixed cosmological time t , the average on the whole particle horizon is:

$$\langle \psi(t) \rangle = f_w(t)\psi_w(t) + f_v(t)\psi_v(t), \quad (16)$$

where the subscripts v and w refer obviously to the values of ψ in voids and walls.

Astrophysical data refer to the local value of ψ that can be generally different from its averaged one $\langle \psi \rangle$. If we consider the universe as a cosmic web, these averaged parameters are physical quantities obtained from an average procedure over a given astrophysical sample containing voids and walls. Moreover, $\psi_w(t)$ and $\psi_v(t)$ are mean values calculated separately in walls and voids.

In any case, the model has a scale of statistical homogeneity, beyond which it has a Friedmann metric evolving with $a(t)$ given by eq.(3). By consequence, the cosmological parameters obtained as averaged quantities at the scale of homogeneity can be straightforward compared with the corresponding values of the standard concordance model. For this reason, we expect that the numerical values for the cosmological parameters of the Λ FB model are similar to those of the standard cosmological model.

As a further consideration on the Λ FB model, note that the formalism introduced allows easily to compute the limit for $\Omega_{k0} \rightarrow 0$ in (8)-(9). At this aim we should just change the hyperbolic metric (2) with parabolic void metric.

$$ds_v^2 = -dt^2 + a_v^2 [d\eta_v^2 + \eta_v^2 d\Omega^2]. \quad (17)$$

This case is an interesting one: if $\Omega_{k0} = 0$ and $\Omega_{m0} + \Omega_{\Lambda0} > 1$, equation (7) implies that $\Omega_Q \leq 0$. The latter means that the flat void region described by metric (17) should be an underdensity. Equation (17) is relevant only to compute the distance-redshift relation, i.e. when considering data on the light cone. Then, we can study equation (15) also in the limit $\Omega_{k0} = 0$. In this case, we get the Λ CDM model by setting $f_{v0} = \epsilon_i = 0$; in another

words, at odds with FB model, the Λ FB contains the Λ CDM model for the aforementioned choice of the parameters.

3 Constraints on the parameter ϵ_i

From equation (5) and (10) we have (for $t = t_i$):

$$\epsilon_i = 1 - \frac{(1 - f_{vi})}{I_{wi}^2 \Omega_{mi}}, \quad I_{wi} = 1 - f_{vi} + f_{vi} \frac{H_{vi}}{H_{wi}} = \frac{H_i}{H_{wi}}. \quad (18)$$

In the original FB model, the parameter I_w is interpreted as a phenomenological lapse function and we have $I_{wi} = 1, \Omega_{mi} \simeq 1, \epsilon_i \ll 1$. By consequence, in the FB model the parameter ϵ_i has no role. On the contrary, in our model (Λ FB), I_w is not a lapse function but simply a measure of the ratio $\frac{H}{H_w}$ or in another words of the ratio between the expansion rate of voids and walls. By consequence, ϵ_i is only constrained to be ≤ 1 , from the existence of II member of eq.(8). It is also possible to express, by means of (15), ϵ_i in terms of the other current cosmological parameters. We have

$$\begin{aligned} \epsilon_i &= \frac{-b \pm 4\Omega_{m0}\sqrt{\Delta}}{2a} \\ b &= -2\Omega_{m0}\Omega_{\Lambda0} + 2f_{v0}\Omega_{m0}\Omega_{k0} + 4f_{v0}\Omega_{m0}\Omega_{\Lambda0} - 4f_{v0}\Omega_{m0} - \\ &\quad -2\Omega_{m0}^2 + 2f_{v0}\Omega_{m0}^2 + 2\Omega_{m0}, \\ a &= \Omega_{m0}^2, \quad \Delta = f_{v0}(\Omega_{k0} + \Omega_{m0} + \Omega_{\Lambda0} - 1)(1 - f_{v0}). \end{aligned} \quad (19)$$

Care must be taken to avoid spurious solutions. In particular, for $f_{v0} \leq 0.1$ we generally have one root, the greater one in (19). The existence of the solution requires $\Omega_{m0} + \Omega_{\Lambda0} + \Omega_{k0} \geq 1$, which implies for equation 7 $\Omega_Q \leq 0$, i.e. a non positive backreaction. This is a check of the consistency of the model, since the partition chosen with voids with an average negative curvature should imply a non positive backreaction, as it is.

4 Constraints on the cosmological parameters of the Λ FB model

4.1 The current curvature Ω_{k0}

In this section we discuss the constraints on the cosmological parameters we can get from the observational data. Equation (15) can be solved with respect to Ω_{k0} and we get:

$$\Omega_{k0} = \frac{\left[1 - \sqrt{(1 - f_{v0})[\Omega_{m0}(1 - \epsilon_i) + \Omega_{\Lambda0}(1 - f_{v0})]}\right]^2}{f_{v0}} - f_{v0}\Omega_{\Lambda0} - \Omega_{m0}\epsilon_i. \quad (20)$$

Current estimates of Ω_{k0} come from CMB and SZ effect data (see [43, 44, 45]). Such measures can be interpreted in the standard framework (Λ CDM) or in the void models (as the FB, the present Λ FB, etc.). In the Λ CDM model, which assumes a constant spatial curvature at fixed time, the observations imply $|\Omega_{k0}| < 0.01$. Remember that within the Buchert scheme, this limit can also be taken into account, provided that it is intended as a mean quantity. Such small value is also consistent with the inflationary scenario. On the contrary in void models, the curvature acts as an effective dark energy. For this reason and in order to fit the observational data, void models require a quite large curvature. Such large value has been claimed consistent to the observation of big voids in the large scale structure. Unfortunately such scenario has two additional implications: a low value for the Hubble constant and/or the existence of a giant void around our location (see [46]). Both of them appear quite implausible. The present model Λ FB on the contrary, since incorporates voids and dark energy, gives the possibility to explore the compatibility between dark energy, large voids, a standard range of values for $\Omega_{m0} \in [0.25, 0.35]$ and a small value for Ω_{k0} in agreement with the WMAP constraint. From the formula (19) we see that a relevant volume void fraction f_{v0} is compatible with a rather small value of Ω_{k0} . For examples considering $\Omega_{\Lambda0} = 0.7$ and $\Omega_{m0} = 0.3$ we have:

- $f_{v0} = 0.7$ for $\Omega_{k0} = 0.0039, \epsilon_i = 0.5$
- $f_{v0} = 0.5$ for $\Omega_{k0} = 0.001, \epsilon_i = 0.16$
- $f_{v0} = 0.2$ for $\Omega_{k0} = 0.01, \epsilon_i = 0.46, \epsilon_i = -0.07$

Note that can exist values for f_{v0} obtained with two values for ϵ_i .

Decreasing $\Omega_{\Lambda0}$ in order to get a similar value for f_{v0} , it requires a larger Ω_{k0} , in agreement with the inhomogeneous models where dark energy is mimicked by the curvature.

Note that for the same range of parameter values, the backreaction term is $\Omega_Q \approx -0.06$ and thus remains relatively small.

Larger values for Ω_Q can be obtained with a small Ω_{k0} and a reasonable void fraction f_{v0} , for a smaller $\Omega_{\Lambda0}$, but a larger Ω_{m0} value. It is worth to stress that our result is not a validation of the standard value for Ω_{k0} . In fact, the current standard constraint $|\Omega_{k0}| \leq 0.01$ is obtained within the Friedmann paradigm. A larger value $\Omega_{k0} \approx 1$ can fit as well the experimental data (CMB and BAO), but it is at odds with inflationary paradigm. Our result is indeed a falsification of a common claim, i.e. large density contrast $|\delta\rho/\rho| \simeq 1$ due to voids necessarily implies a large curvature contrast $|\delta\mathcal{R}/\mathcal{R}| \simeq 1$. In our model a large fraction of voids is compatible with a small spatial curvature and a standard value for cosmological constant.

4.2 Current volume voids fraction f_{v0}

An interesting feature of the Λ FB model is the constraints on the present day volume void fraction f_{v0} from the values of the other cosmological parameters. By posing the condition $\Omega_{k0} \geq 0$, $\Omega_{m0} > -2\Omega_{\Lambda0} + 2\sqrt{\Omega_{\Lambda0}}$ (which is always satisfied) and neglecting spurious solutions we have

$$\begin{aligned} f_{v0} &\geq X_0, \quad X_0 = \frac{-b + 4\sqrt{\Delta}}{2a}, \\ a &= 4\Omega_{\Lambda0}^2 - 4\Omega_{\Lambda0} + \Omega_{m0}^2 + 4\Omega_{m0}\Omega_{\Lambda0}, \\ \Delta &= (\Omega_{m0} + \Omega_{\Lambda0} - 1) [\Omega_{\Lambda0}(\Omega_{m0} + \Omega_{\Lambda0} - 1) + \epsilon_i \Omega_{m0}^2 (1 - \epsilon_i)], \\ b &= -4\Omega_{m0}\epsilon_i + 4\Omega_{m0}\Omega_{\Lambda0}\epsilon_i + 2\Omega_{m0}^2\epsilon_i - 2\Omega_{m0}^2 - 6\Omega_{m0}\Omega_{\Lambda0} + 4\Omega_{\Lambda0} + \\ &\quad + 2\Omega_{m0} - 4\Omega_{\Lambda0}^2. \end{aligned} \tag{21}$$

Note that, if $\Omega_{m0} + \Omega_{\Lambda0} < 1$, no limitations arise for the actual volume void fraction f_{v0} . For $\Omega_{m0} + \Omega_{\Lambda0} = 1$ we obtain $X_0 = \epsilon_i$.

Finally, note that in the case $\Omega_{k0} = 0$ (for all times), the inequality (21) becomes an equality. As a result, in this case we can obtain the standard concordance model ($f_{v0} = 0$) or a model with voids with a flat Friedmann metrics and $f_{v0} = X_0$, provided that $\Omega_{m0} + \Omega_{\Lambda0} > 1$.

We study now the equation (15) in terms of the allowed current volume voids fraction f_{v0} . The equation can be solved in terms of f_{v0} with some little algebra. Care must be taken for the possible appearance of spurious solutions, by solving it graphically (i.e. equation (15)) and eliminating the latter ones. Then expliciting equation (15) in terms of f_{v0} we get:

$$\begin{aligned} f_{v0} &= \frac{-b \pm 4\sqrt{\Delta}}{2a}, \\ a &= 4\Omega_{\Lambda0}^2 + \Omega_{m0}^2 + \Omega_{k0}^2 - 4\Omega_{\Lambda0} + 4\Omega_{\Lambda0}\Omega_{k0} + 2\Omega_{m0}\Omega_{k0} + 4\Omega_{m0}\Omega_{\Lambda0}, \\ b &= 4\Omega_{\Lambda0} + 2\Omega_{m0} - 6\Omega_{\Lambda0}\Omega_{m0} - 2\Omega_{k0} - 2\Omega_{\Lambda0}\Omega_{k0} - 2\Omega_{m0}^2 - 4\Omega_{\Lambda0}^2 - \\ &\quad - 2\Omega_{k0}\Omega_{m0} + 4\Omega_{m0}\Omega_{\Lambda0}\epsilon_i + 2\Omega_{m0}^2\epsilon_i + 2\Omega_{k0}\Omega_{m0}\epsilon_i - 4\Omega_{m0}\epsilon_i, \\ \Delta &= (\Omega_{\Lambda0} + \Omega_{m0} + \Omega_{k0} - 1)[\Omega_{m0}\Omega_{\Lambda0} + \Omega_{\Lambda0}^2 + \Omega_{\Lambda0}\Omega_{k0} - \Omega_{\Lambda0} + \\ &\quad + \Omega_{m0}^2\epsilon_i(1 - \epsilon_i) + \Omega_{m0}\Omega_{k0}(1 - \epsilon_i)]. \end{aligned} \tag{22}$$

Note that for $\Omega_{\Lambda0} + \Omega_{m0} > 1$ at least a solution is always present, while for $\Omega_{\Lambda0} + \Omega_{m0} < 1$ we can have no solutions. In any case for $\Omega_{m0} + \Omega_{\Lambda0} + \Omega_{k0} < 1$, since $\Omega_Q < 0$, according to equation (7), no solutions are available. For an example of possible values, see the table (1). Quite generally, we can have two possible values for f_{v0} :

- $f_{v01} < 1$
- $f_{v02} \approx 0.1$ or greater

This implies that if the initial fraction of voids is:

$\Omega_{\Lambda 0}$	Ω_{m0}	Ω_{k0}	ϵ_i	f_{v01}	f_{v02}
0.7	0.28	0.01	0	<i>no</i>	<i>no</i>
0.7	0.28	0.02	0	<i>no</i>	0.066
0.7	0.28	0.021	0	0.03	0.14
0.7	0.28	0.022	0	0.023	0.18
0.7	0.29	0.018	0	0.0026	0.32
0.7	0.29	0.018	0.01	0.004	0.33
0.7	0.30	0.001	0	<i>no</i>	0.043
0.7	0.30	0.01	0	<i>no</i>	0.32
0.7	0.30	0.02	0	<i>no</i>	0.51
0.7	0.30	0.01	0.01	0.0002	0.34
0.7	0.30	0.01	0.1	0.016	0.44
0.7	0.305	0.0007	0	<i>no</i>	0.18
0.7	0.305	0.01	0	<i>no</i>	0.4
0.7	0.31	0.01	0	<i>no</i>	0.47
0.7	0.31	0.01	0.01	<i>no</i>	0.48
0.7	0.31	0.01	0.15	0.014	0.62

Table 1: Allowed actual volume voids fraction for Λ FB model.

- $f_{vi} < f_{v01}$ then at $t = t_0$ $f_{v0} = f_{v01}$
- $f_{vi} > f_{v01}$ then at $t = t_0$ $f_{v0} = f_{v02}$

Considering the value for f_{vi} estimated from WMAP data ($\approx \in [10^{-5}, 10^{-2}]$) and the values of f_{v01} and f_{v02} , both of the previous cases could be fulfilled. According to the model, starting from a fraction of voids f_{vi} within WMAP constraints, we could get at the present time t_0 both f_{v01} and f_{v02} . Which one is a question of the precise value of f_{vi} with respect to the f_{v01} . The dependence from the other parameters is quite complex. In fact from the table (1) is apparent that for a choice $\Omega_{\Lambda 0} \simeq 0.7$, $\Omega_{k0} \leq 0.02$ and $0 \leq \epsilon_i \leq 0.15$, changing Ω_{m0} from 0.28 to 0.31 produces a variation of f_{v0} from ~ 0.0002 up to $\simeq 0.6$. In the Λ FB, contrary to the FB one, the final fraction of voids f_{v0} is dependent from the initial one f_{vi} . This happens because in the Λ FB model no tracker solutions are available.

4.3 Current cosmological constant $\Omega_{\Lambda 0}$

As mentioned, both FB and Λ FB models collapsing regions are within walls; for this reason the model has $\Omega_{k0} \geq 0$ and we get:

$$\Omega_{\Lambda 0} \leq G \quad (23)$$

$$G = \frac{\epsilon_i \Omega_{m0} - 2f_{v0} \epsilon_i \Omega_{m0} + 1 - 2\Omega_{m0} f_{v0}^2 - 2f_{v0} + 3f_{v0} \Omega_{m0} - \Omega_{m0} + 2f_{v0}^2 - 2\sqrt{Q}}{(1 - 2f_{v0})^2},$$

$$Q = f_{v0}(1 - f_{v0})[2f_{v0}^2\Omega_{m0} - f_{v0}^2 + f_{v0} - f_{v0}\Omega_{m0} + \epsilon_i\Omega_{m0}(1 - 2f_{v0})].$$

Note that $Q > 0$ for all values allowed of f_{v0}, Ω_{m0} . We conclude this subsections by analysing the formula (15) from the point of view of $\Omega_{\Lambda0}$. This is in our opinion the main point of the present study, since it allows to estimate the fraction of $\Omega_{\Lambda0}$ due to the inhomogeneities in the large scale structure. By solving (15) with respect to $\Omega_{\Lambda0}$ we obtain

$$\begin{aligned} \Omega_{\Lambda0} &= \frac{-b - 4\sqrt{f_{v0}(1 - f_{v0})\sqrt{\Delta}}}{2a}, \\ a &= 1 + 4f_{v0}^2 - 4f_{v0}, \\ b &= -6\Omega_{m0}f_{v0} + 4f_{v0} - 2 + 4f_{v0}\Omega_{m0}\epsilon_i - 4f_{v0}^2 + 2\Omega_{m0} + 4f_{v0}^2\Omega_{m0} + \\ &\quad + 4f_{v0}^2\Omega_{k0} - 2f_{v0}\Omega_{k0} - 2\Omega_{m0}\epsilon_i, \\ \Delta &= 2f_{v0}^2\Omega_{m0} - f_{v0}^2 + f_{v0} - f_{v0}\Omega_{m0} + \epsilon_i\Omega_{m0}(1 - 2f_{v0}) + 2f_{v0}^2\Omega_{k0} - \\ &\quad - 3f_{v0}\Omega_{k0} + \Omega_{k0}. \end{aligned} \tag{24}$$

To our knowledge this is the first equation which allows to compute the behaviour of $\Omega_{\Lambda0}$ versus in particular f_{v0} without imposing simple assumptions (as spherical or special symmetries) on inhomogeneities. First of all, from WMAP data we put a maximum allowed mean value for $\Omega_{k0} \simeq 0.01$. For Ω_{m0} we consider range of values (0.2, 0.35) (see for example [47]). In fig 1, 2, 3 we plot $\Omega_{\Lambda0}$ vs. f_{v0} and Ω_{m0} with $\Omega_{k0} = 0.01$ and $\{\epsilon_i\} = \{0, 0.1, 0.9\}$. As




Figure 1: $\Omega_{\Lambda0}$ vs. Ω_{m0} and f_{v0} . $\epsilon_i = 0$. The labelled curves are the isocurves for the corresponding value of $\Omega_{\Lambda0}$.

apparent from the figures, a set of standard Λ CDM values for $\Omega_{\Lambda0}, \Omega_{m0}, \Omega_{k0}$




Figure 2: $\Omega_{\Lambda 0}$ vs. Ω_{m0} and f_{v0} . $\epsilon_i = 0.1$. For the isocurves see caption fig1




Figure 3: $\Omega_{\Lambda 0}$ vs. Ω_{m0} and f_{v0} . $\epsilon_i = 0.9$. For the isocurves see caption fig1

is compatible with a quite large fraction of voids (roughly up to 0.8). In particular, we see that the maximum value $\Omega_{m0} \simeq 0.35$ together with the maximum allowed mean value for Ω_{k0} predicted by WMAP ($\simeq 0.01$) is compatible with the minimum value $\Omega_{\Lambda 0} \simeq 0.65$ and a large fraction of voids (up to $\simeq 0.8$). If this were the case, spatial inhomogeneities could account

for a maximum percentage $\approx 10\%$ with respect to the concordance value $\Omega_{\Lambda 0} \simeq 0.73$. Moreover, note that for $\epsilon = 0$ or ~ 0.1 we have that huge values for $\Omega_{\Lambda 0} > 0.8$ can be obtained with $\Omega_{m0} \in [0.25, 0.35]$ together with a huge value for $f_{v0} > 0.8$. Conversely, for high $\epsilon \sim 0.9$ we have that with $\Omega_{m0} \in [0.25, 0.35]$, $\Omega_{\Lambda 0} > \simeq 0.73$ can be obtained for $f_{v0} < \simeq 0.1$. This can help to understand the interesting role of ϵ in the Λ FB model. Finally, we consider the case with $\epsilon = \Omega_{k0} = 0$ where a situation close to figure 1 arises. Note that the concordance model is regained by further setting $f_{v0} = 0$. As explained at the end of section 2, this case corresponds to the physical situation where both walls and voids have mean zero spatial curvature at any time but where voids represent an underdensity. Hence, voids expand faster than walls and as a result a non trivial volume void fraction $f_v(t)$ with $f_{v0} \neq 0$ emerges as the case with $\Omega_{k0} \neq 0$.

5 Age of the universe compatible with the Λ FB model

An important test in cosmology is given by the age of the universe. The age of globular cluster implies the universe should be certainly older than $> 12\text{Gyr}$. In this section we give an estimation of the age of the universe predicted by Λ FB model. First of all, from equation (13) evaluated at present time t_0 we get:

$$t_0 = \frac{2}{3H_0\sqrt{\Omega_{\Lambda 0}}} \ln \left(\sqrt{(1-f_{v0})\frac{\Omega_{\Lambda 0}}{\Omega_{m0}(1-\epsilon_i)}} + \sqrt{(1-f_{v0})\frac{\Omega_{\Lambda 0}}{\Omega_{m0}(1-\epsilon_i)} + 1} \right) \quad (25)$$

where H_0 is the present cosmological constant at the scale of homogeneity. A consistent value for t_0 ($> 12\text{Gyr}$) is always possible with a large volume fraction of voids.

Let's give an example: let us consider $t_0 \geq 13\text{ Gyr}$, a restrictive value for $\epsilon_i = 0$, a large amount of dark energy ($\Omega_{\Lambda 0} \in [0.67, 0.75]$) and a relatively small curvature, i.e. $\Omega_{k0} \leq 0.08$. For H_0 we consider the range $(55-75)\text{km/s/Mpc}$, since a smaller value appears in contrast with the actual data. From eq.(25) we get a reasonable t_0 with a void fraction $f_{v0} \approx (40-50)\%$ and $H_0 \approx (55-60)\text{km/s/Mpc}$. Increasing H_0 up to $\simeq 70\text{km/s/Mpc}$ and considering $\epsilon_i < 1$, we got an age $t_0 \geq 13\text{Gyr}$ with $f_{v0} \approx \in (0, 0.3)$. This shows how the model can account for a reasonable choice of the cosmological parameters (age, H_0 , curvature, etc..) with a fraction of voids as large as 30%. An even greater fraction of voids, $f_{v0} > 0.3$, and $H_0 \simeq 70\text{km/s/Mpc}$ require a larger value of ϵ_i and more precisely $\epsilon_i \sim f_{v0}$.

$\Omega_{\Lambda 0}$	Ω_{m0}	f_{v0}	H_0	t_0
0.67	0.26	0.4	55	15.1
0.67	0.26	0.4	60	13.9
0.67	0.26	0.5	55	14.1
0.67	0.31	0.4	55	14.2
0.67	0.31	0.4	60	13.0
0.67	0.31	0.5	55	13.2
0.67	0.34	0.4	55	13.6
0.70	0.27	0.4	55	14.9
0.70	0.27	0.4	60	13.6
0.70	0.27	0.5	55	13.9
0.70	0.30	0.4	55	14.3
0.70	0.30	0.4	60	13.1
0.70	0.30	0.5	55	13.3
0.71	0.26	0.4	55	15.0
0.71	0.26	0.4	60	13.8
0.71	0.26	0.5	55	14.0
0.73	0.26	0.4	60	13.7
0.73	0.26	0.5	55	14.0
0.75	0.26	0.4	60	13.7

Table 2: Age of the universe compatible with AFB model.

6 Dressing cosmological parameters and the distance angular function

One of the issue of the Buchert averaging scheme is how to relate volume average quantities which are non-local to observable ones, which are local. In contrast to many approaches to the Buchert equations which usually neglect this fundamental issue and following Wiltshire [23, 41], we match the radial null section of the wall metric (1), rewritten as

$$\begin{aligned}
ds_w^2 &= -dt^2 + a_w^2 [d\eta_w^2 + \eta_w^2 d\Omega^2] \\
&= -dt^2 + \frac{(1-f_v)^{\frac{2}{3}} a^2}{f_{wi}^{\frac{2}{3}}} [d\eta_w^2 + \eta_w^2 d\Omega^2].
\end{aligned} \tag{26}$$

with the metric at the scale of homogeneity given by

$$ds^2 = -dt^2 + a^2 d\eta^2 + A(t, \eta) d\Omega^2, \tag{27}$$

$A(t, \eta)$ is an area function satisfying $4\pi \int_0^{\eta_{\mathcal{H}}} A d\eta = a^2 V_i(\eta_{\mathcal{H}})$, where $\eta_{\mathcal{H}}$ is the particle horizon radius. Practically, the wall observer must dress the cosmological parameter and not simply to relate the volume average scale

factor to the observed redshift z . The dressed wall geometry is:

$$ds_w^2 = -dt^2 + a^2 \left[d\eta^2 + (1 - f_v)^{\frac{2}{3}} f_{wi}^{-\frac{2}{3}} \eta_w^2 d\Omega^2 \right], \quad (28)$$

with

$$d\eta_w = \frac{f_{wi}^{\frac{1}{3}} d\eta}{(1 - f_v)^{\frac{1}{3}}}, \quad \eta = \int_t^{t_0} \frac{dt}{a}. \quad (29)$$

It is by means of the metric (28) that the wall observer, in galaxies, measures the distance-reashift function $d_L(z)$. The angular-distance relation $d_A(t)$ (remember that $d_L = (1 + z)^2 d_A$) is:

$$d_A(z) = \frac{a_0}{(1 + z)} \bar{\eta}_w, \quad 1 + z = \frac{a_0}{a}, \quad (30)$$

$$\bar{\eta}_w = (1 - f_v)^{\frac{1}{3}} \int_t^{t_0} \frac{dt}{(1 - f_v)^{\frac{1}{3}} a}.$$

Using eq.(13), expression (30) becomes:

$$d_A(t) = \sinh^{\frac{2}{3}} \left(\frac{3}{2} H_0 \sqrt{\Omega_{\Lambda 0} t} \right) \int_t^{t_0} \frac{dt}{\sinh^{\frac{2}{3}} \left(\frac{3}{2} H_0 \sqrt{\Omega_{\Lambda 0} t} \right)}, \quad (31)$$

where t_0 is given by eq.(25). We can see that d_A has the same functional form versus the time t both for Λ CDM and Λ FB models. The only difference is the expression of the function t_0 . Obviously the expression for d_A changes with respect to the concordance model when expressed in terms of the redshift z . To express $t(z)$ along the past null cone, we must to know the function $a(t)$ which, for the Λ FB model can be obtained by integrating numerically the equations (9).

7 Conclusions

We have presented a preliminary study of the recent Λ FB cosmological model. An explicit formula relating all the cosmological parameters of the Λ FB model is obtained and analysed. The relevant feature of the Λ FB model is the presence of the cosmological constant and spatial inhomogeneities without spherical symmetry [39]. By consequence, the standard Λ CDM model can be recovered with a suitable choice of the parameters. The model allows to analyse the departures from the standard cosmological model without invoking perturbation theory.

The first result of the present study is the consistency of a large volume voids fraction (> 0.1) with a small spatial curvature, even within WMAP constraint ($|\Omega_{k0}| \leq 0.01$). This falsifies the argument often used in the literature to rule out dark energy, i.e. a large fraction of voids observed

necessarily implies a large negative value for the curvature .

The second result is the important role (absent in the FB model) of the initial volume void fraction f_{vi} . In fact, for a reasonable range of values for the cosmological parameters $\Omega_{m0}, \Omega_{k0}, \Omega_{\Lambda0}$ and dependent on the parameter ϵ_i , the model generally provides two possible values for the current volume void fraction: $f_{v01} \ll 1$ and $f_{v02} \geq 0.1$. If the initial value $f_{vi} < f_{v01}$, then the universe evolves up to a final volume void fraction f_{v01} , while for $f_{vi} > f_{v01}$ the system evolves up to the second root f_{v02} . As it is evident from the table (1), the model is quite sensitive to small variations of cosmological parameters. Finally, we analysed the formula (24) giving the exact value for $\Omega_{\Lambda0}$ in terms of the other current cosmological parameters. Setting the maximum value allowed for $\Omega_{k0} \simeq 0.01$ and $\Omega_{m0} \simeq 0.35$, we get (see figures 1, 2, 3) that the lower value for $\Omega_{\Lambda0} \simeq 0.65$ is compatible with a volume void fraction $\in (0, 0.8)$. Hence, considering a total amount of dark energy 73% predicted by the Λ CDM model, the inhomogeneities could account for a maximum percentage $\approx 10\%$ of $\Omega_{\Lambda0}$.

As a final consideration, note that also in the Λ FB model it is possible to add clock effects present in [23]. This can be done simply by considering I_w as a lapse function together with $\epsilon \ll 1 \sim f_{vi}$ (see [41]). Equations (8), (9) and formula (15) are left unchanged. The changes are in the dressing procedure, i.e. equations (26)-(31). Moreover, in the case of non vanishing lapse function the cosmological parameters must be 'dressed' [22, 23]. As an example, if clock effects are present, Ω_m is the bare volume-average density parameter while the measured density parameter in walls Ω_{mw} is given by $\Omega_{mw} = I_w^3 \Omega_m$. This is a preliminary parametric study on the relation between cosmological parameters at the present time. In a future paper we intend to study the complex problem to fit the observational data by integrating numerically the model field equations.

Acknowledgements

We would like to thank D. L. Wiltshire for useful comments and suggestions.

References

- [1] B.S. Perlmutter, et al., *Astrophys. J.* **483** (1997) 565.
- [2] A.G. Riess, et al., *Astron. J.* **116** (1998) 1009.
- [3] B.S. Perlmutter, et al., *Astrophys. J.* **517** (1999) 565.
- [4] M.N. C  lerier, *Astron. and Astrophys.* **353** (2000) 63.
- [5] M. Tanimoto and Y. Nambu, *Class. Quantum Grav.* **24** (2007) 3843.

- [6] J.W. Moffat, *JCAP* **0605** (2006) 001.
- [7] K. Enqvist, *Gen. Relativ. Gravit.* **40** (2008) 451.
- [8] H. Alnes, M. Amarzguioui, and O. Gron, *Phys. Rev. D* **73** (2006) 083519.
- [9] W. Valkenburg, *JCAP* **0906**(2009) 010.
- [10] T. Biswas, A. Notari and W. Valkenburg, *JCAP* **1011**(2010) 030
- [11] W. Valkenburg, *JCAP* **1201**(2012) 047.
- [12] K. Tomita, *Prog. Theor. Phys.* **106** (2001) 929.
- [13] A. Paranjape, and T.P. Singh, *Class. Quantum Grav.* **23** (2006) 6955.
- [14] V. Marra, E.W. Kolb, S. Matarrese, and A. Riotto, *Phys. Rev. D* **76** (2007) 123004.
- [15] T. Buchert, and M. Carfora, *Class. Quantum Grav.* **19** (2008) 195001.
- [16] V. Marra, E.W. Kolb, and S. Matarrese, *Phys. Rev. D* **77** (2008) 023003.
- [17] S. Räsänen, *JCAP* **0611** (2006) 003.
- [18] T. Buchert, *Gen. Relativ. Gravit.* **32** (2000) 105.
- [19] T. Buchert, *Gen. Relativ. Gravit.* **40** (2008) 467.
- [20] T. Buchert, and M. Carfora, *Phys. Rev. Lett.* **90** (2003) 031101.
- [21] T. Buchert, and M. Carfora, *Class. Quantum Grav.* **19** (2002) 6109.
- [22] D.L. Wiltshire, *Phys. Rev. Lett.* **99** (2007) 251101.
- [23] D.L. Wiltshire, *New J. Phys.* **9** (2007) 377.
- [24] J. Kwan, M. J. Francis and G. F. Lewis, *Mon. Not. R. Astron. Soc. Lett.* **399** (2009)L6.
- [25] P.R. Smale and D.L. Wiltshire, *Mon. Not. R. Astron. Soc.* **413** (2011) 367.
- [26] R.A. Vanderveld, É.É. Flanagan, and I. Wasserman, *Phys. Rev. D* **74** (2006) 023506.
- [27] A.E. Romano, *JCAP* **05** (2010) 020.
- [28] A. Krasinski, C. Hellaby, K. Bolejko, and M.N. Célérier, *General Relativ. Gravit.* **42** (2010) 2453.

- [29] M.N. C  lerier, K. Bolejko, and A. Krasinski, *Astron. Astrophys* **A21** (2010) 518.
- [30] P.M.Sutter, et al., *Astrophys. J.* **761**, 44 (2012).
- [31] F. Hoyle and M.S. Vogeley, *Ap. J.* **607** (2004) 751.
- [32] D.C. Pan, M.S. Vogeley, F. Holey, Y-y Choi and C. Park, arXiv:1103.4156.
- [33] A.V. Tikhonov and I.D. Karachentsev, *Astrophys. J.* **653** (2006) 369.
- [34] M. Zaldarriaga, D.N. Spergel and U. Seljak, *Astrophys. J.* **488** (1997) 1.
- [35] E. Komatsu et al., *Astrophys. J.S.* **180** (2009) 330.
- [36] J.M.Virey at al., *JCAP* **12** (2008).
- [37] L. Amendola, K. Kainulainen, V. Marra and V. Quartin, *Phys. Rev. Lett.* **105** (2010) 121302.
- [38] V. Marra and M. Paakkonen, *JCAP***12** (2010) 021.
- [39] A.E. Romano and C. Pisin*JCAP* **016**, (2011) 1110.
- [40] V. Marra and A. Notari, *Class. Quant. Grav.* **28** (2011)164004.
- [41] S. Viaggiu, *Class. Quantum Grav.* **29** (2012) 035016.
- [42] S. Viaggiu, *Class. Quantum Grav.* **27** (2010) 155002.
- [43] J.N. Dossett and M. Ishak, arXiv:1205.2422.
- [44] A.H Guth and Y. Nomura, arXiv:1203.6876.
- [45] Y. Wang and P. Mukherjee, *Phys. Rev. D* **76** (2007) 103533.
- [46] T. Clifton, P.G. Ferreira and J. Zuntz, *JCAP* **07** (2009) 029.
- [47] J. A. Peacock et al., *Nature* **410** (2001) 169.